

REVIEW

Open Access



# Short-chain fatty acids as a link between diet and cardiometabolic risk: a narrative review

Eline Birkeland<sup>1</sup>, Sedegheh Gharagozian<sup>1</sup>, Jørgen Valeur<sup>2</sup> and Anne-Marie Aas<sup>1,3\*</sup>

## Abstract

**Aim** Diet has a profound impact on cardiometabolic health outcomes such as obesity, blood glucose, blood lipids and blood pressure. In recent years, the gut microbiota has emerged as one of several potential key players explaining dietary effects on these outcomes. In this review we aim to summarise current knowledge of interaction between diet and gut microbiota focusing on the gut-derived microbial metabolites short-chain fatty acids and their role in modulating cardiometabolic risk.

**Findings** Many observational and interventional studies in humans have found that diets rich in fibre or supplemented with prebiotic fibres have a favourable effect on the gut microbiota composition, with increased diversity accompanied by enhancement in short-chain fatty acids and bacteria producing them. High-fat diets, particularly diets high in saturated fatty acids, have shown the opposite effect. Several recent studies indicate that the gut microbiota modulates metabolic responses to diet in, e.g., postprandial blood glucose and blood lipid levels. However, the metabolic responses to dietary interventions, seem to vary depending on individual traits such as age, sex, ethnicity, and existing gut microbiota, as well as genetics. Studies mainly in animal models and cell lines have shown possible pathways through which short-chain fatty acids may mediate these dietary effects on metabolic regulation. Human intervention studies appear to support the favourable effect of short-chain fatty acid in animal studies, but the effects may be modest and vary depending on which cofactors were taken into consideration.

**Conclusion** This is an expanding and active field of research that in the near future is likely to broaden our understanding of the role of the gut microbiota and short-chain fatty acids in modulating metabolic responses to diet. Nevertheless, the findings so far seem to support current dietary guidelines encouraging the intake of fibre rich plant-based foods and discouraging the intake of animal foods rich in saturated fatty acids.

**Keywords** Diet, Short-chain fatty acids, Gut microbiota, Human studies, Metabolic effects

## Introduction

The role of diet in health and chronic conditions such as obesity, insulin resistance, and cardiovascular disease is well known [1] and recognized in clinical guidelines [2, 3].

Diet also shapes the composition of the gut microbiota, which in recent years has emerged as one of several potential key players explaining dietary effects on health and disease [4, 5]. However, studying the relationship between health and microbiota in humans is difficult due to challenges of controlling for environmental factors in study subjects. Lately, a number of large-scale studies including more than 800 people have identified gut microbiota-diet interactions that associate with different cardiometabolic markers [5–7], but so far, only animal studies offer some evidence of causality. The human studies also reveal that the metabolic responses to food varies substantially partly

\*Correspondence:

Anne-Marie Aas  
a.m.aas@medisin.uio.no

<sup>1</sup> Section of Nutrition and Dietetics, Department of Clinical Service, Division of Medicine, Oslo University Hospital, Oslo, Norway

<sup>2</sup> Unger-Vetlesen Institute, Lovisenberg Diaconal Hospital, Oslo, Norway

<sup>3</sup> Institute of Clinical Medicine, University of Oslo, Oslo, Norway



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

due to individual differences in gut microbial composition and functions. Numerous studies also report that the gut microbiome of people with diseases, such as type 2 diabetes, stroke and immune-mediated inflammatory disease, is distinctly different from that of healthy individuals as they present with a microbiome with less diversity and reduced abundance of health promoting species [8–13].

Unravelling the interactions between diet and the gut microbiota and their impact on metabolism and cardiometabolic disease could open for new approaches to obtain good health and prevent and treat disease by feeding our gut microbiota the optimal diet. What the best diet is may differ from one individual to another depending on metabolic phenotype, existing microbiome and more [7, 14]. Identifying predictors of metabolic responses is another research field that needs to be mapped and tested in intervention trials. Finally, such research can be useful in the development of dietary guidelines and help optimize personalized diet recommendations based on prediction models derived from large studies of diet-microbiota interactions and effect on cardiometabolic health.

In this article, we review the current knowledge of diet and gut microbiota interaction focusing on the gut-derived microbial metabolites of short-chain fatty acids (SCFA) and their role in modulating cardiometabolic disease risk. Other dietary derived compounds produced by bacteria, such as trimethylamine (TMA), other methylamines, polyamines, and secondary bile acids may also affect host health, but is outside the scope of this narrative review.

**Method**

In this narrative review, we did not perform a systematic search for articles, but publications were selected on a discretionary basis for their relevance to the aim of the review. We focused on studies in humans but included some studies in animal models and cell lines in cases where we found it relevant to underpin possible pathways of action, where these studies guided and supported findings in human studies and in the cases where human studies were lacking. Most included articles used were chosen from searches in Pubmed and Google scholar from May 2021 to July 2022 using the keywords: diet, short-chain fatty acids, gut microbiota, gut bacteria,

<b>Phylum</b>	Firmicutes
<b>Family</b>	Ruminococcaceae
<b>Genus</b>	<i>Faecalibacterium</i>
<b>Species</b>	<i>Faecalibacterium prausnitzii</i>


**Fig. 2** Simplified rank-based classification of the butyrate producer *Faecalibacterium prausnitzii*

cardiometabolic, prebiotics, SCFA receptors, LPS, inflammation, and combinations thereof. Additional publications were added using the snowballing method as a complementary approach, including relevant papers that were cited in already included publications and articles that came to our attention mentioned by colleagues or cited during a conference.

**Gut microbiota**

In clinical science microbiota as a term often describes bacteria. Although the bacteria account for the main mass, the microbiota comprises archaea, protists, fungi and viruses as well (Fig. 1) [15]. The bacteria are ranked into phylum, order, class, family, genus and species (Fig. 2) [16]. The major bulk of bacteria in the gut are the phyla Firmicutes and Bacteroidetes (90%), followed by Actinobacteria and Proteobacteria (Fig. 3) [16]. Cyanobacteria, Verrumicrobia, Tenericutes and other phyla not yet assigned, are included as well [16].

In 2004, a connection between gut microbiota and development of excessive body fat and insulin resistance was discovered in mice [17]. Further research on mice has reported increased capacity to extract energy from undigested food components and that obesity is transmissible between individuals through gut bacteria [18]. This was followed by a number of epidemiologic studies reporting differences in gut bacteria between healthy humans and humans with increased risk for cardiometabolic diseases [8, 9, 12, 19]. These differences include reduced microbial diversity, changed ratio of the two major phyla Firmicutes and Bacteroidetes, reduced concentrations of some species that are assumed healthy, and increased concentrations of others that are considered harmful [8, 9, 12, 19].



**Fig. 1** Overview of microbiota

These differences are often called dysbiosis. Dysbiosis as a term, however, is disputable since it is unclear what constitutes a healthy core microbiota or whether it even exists [20]. So far, speculations about which traits frame a healthy bacterial composition in the gut, are deduced from the corresponding opposite features found between microbiota in healthy people and people with overweight and health problems. Nonetheless, these findings caused a spiked interest in gut microbiota as a future objective for management of cardiometabolic diseases.

### The size of the gut microbiota pool and changing perception of its role

Long ago, the quantity of bacterial cells in the human gut was estimated to largely exceed the number of human cells [21]. A recent recalculation, though, reduced the bacterial cell count to match the number of human cells [21]. Nevertheless, human beings have approximately 22 000 genes, whereas our bacteria all together have a hundred times more, with a far greater genetic capacity to produce and express biologically active compounds.

Our conception of the gut bacteria's impact has altered from docile dwellers in the gut to an active bacterial society, with means to modulate the biology of their human hosts. This host–microbe relationship is believed to benefit human health, because the gut microbiota extracts nutrients from undigested dietary components, maintains the intestinal barrier, protects the host against harmful bacteria, produces essential vitamins, and modulates the immune system [22]. To what extent our gut inhabitants, with their outnumbering genes, are able to regulate human bodily functions, is a hot research field.

### What shapes the gut microbiota?


The bacterial composition changes throughout life and is affected by genetics, age, transit time through the gut, and a wide range of environmental factors, including mechanism of birth, breastfeeding, maternal microbiota, diet, lifestyle, medication, and state of health [22]. Knowledge about which factors that affect gut microbiota has increased considerably during the last years including factors that may have confounded earlier studies. The importance of controlling for diet, geographical

residence, socio-economic status and medication when studying relations between microbiota and cardiometabolic diseases has since been emphasized by several research groups [23–25]. Indeed, environmental factors appear to have even greater impact on the gut microbiota than genetics, and a recent study [26] found that as much as 20% of the variability between microbiota in people was associated with diet, medication and body composition [26]. The same authors estimated the overall microbiome heritability to be between 1.9% and 8.1%. Furthermore, the transmissibility of obesity between mice has also been shown to be diet-dependent and that a diet rich in fruits and vegetables and low in saturated fat appears to be protective [27].

### Gut microbiota composition, diversity, and function

Although the bacterial phyla stay relatively fixed in the healthy adult gut, the species themselves are highly susceptible [28]. Yet, the bacterial community is resilient, i.e., normally re-establishes after temporary disruptions of diet or medication [28]. The microbial diversity can be defined as “the number and abundance distribution of distinct types of organisms” and high diversity is associated with good health [29]. The diversity has been reported to increase during childhood, remain stable in adulthood and decline in old age [30], although a study by Odumaki et al. suggests that the observed decline in elderly people may be confounded by external factors such as residing in own home or in an institution [31].

The gut bacteria have various capabilities and perform different tasks in the colon [29, 32, 33], but different species may possess common abilities. This means that healthy individuals may have dissimilar compositions of well-functioning bacterial communities. Even if the composition of gut microbiota varies considerably between people at species level, the bacterial functions appear to vary less between people than the actual species [29, 33]. A healthy core of microbial functions may thus turn out to be more relevant than a healthy core microbiota composition.



**Fig. 3** The major bacterial phyla in the gut

### Dietary fibres are fermented by the gut microbiota to SCFA

Carbohydrates that humans are unable to digest themselves are referred to as dietary fibres. They pass through the small intestine into the colon where they are fermented by the microbiota [34]. The indigestible but fermentable dietary fibres from plant-based food are the preferred energy source for the gut bacteria [35], which ferment the fibres into various compounds, including SCFA [35].

The fermentation process of fibres from plant-based food increases the luminal acidity in the colon, which provides an environment more suited to healthy bacteria than harmful bacteria [36]. The gut bacteria are also capable of fermenting proteins, but appear to do so only if the fibres from plant-based food are short in supply [36].

Prebiotic fibres are defined as; substrates selectively utilised by host microorganisms conferring a health benefit [37]. During the recent years they have been highlighted as a possible treatment approach in overweight and cardiometabolic diseases [38–43]. Galacto-oligosaccharides (GOS) and inulin-type fructans (ITF) are among the most investigated prebiotics. Clinical trials report beneficial effects on glycaemic regulation, suppression of energy intake and appetite as well as weight loss after treatment with prebiotic fibres, including ITF and GOS [37, 38, 41–46].

### Does the microbiota modulate postprandial responses to diet?

Diet plays a major role in shaping the gut microbiota, but does the gut microbiota modulate the postprandial responses to food intake, and can this explain the heterogeneity in metabolic effects of dietary interventions? In 2015, Zeevi et al. [7] monitored glucose levels continuously for a week in 800 healthy and prediabetic individuals, and measured responses to over 46 000 meals. They found that the response to identical meals were highly variable. A machine-learning algorithm was made based on the measured blood parameters, dietary habits, anthropometrics, physical activity, and gut microbiota in this cohort. The algorithm accurately predicted individual postprandial glycaemic responses to typical meals eaten during the intervention. The authors validated these predictions in a new cohort and performed a randomised controlled trial using this algorithm. A dietary intervention based on the predictions lowered postprandial responses significantly and resulted in consistent changes in gut microbiota composition [7].

More recently, the PREDICT study, with over 1000 men and women, showed that individual factors, like the gut-microbiota, had more impact than macronutrient composition on postprandial lipidemia (7,1% of variance

compared with 3,6%), but not postprandial glycaemia (6% vs 15,4%) [14]. The authors validated their findings in an independent cohort and devised a machine-learning model that predicted both triglyceride and glycaemic response to food intake. Interestingly, genetic variations only partially influenced the predictions (9.5% for glucose, 0.8% for triglycerides, and 0.2% for C-peptide) [14]. In the PREDICT study, deep metagenomics sequencing of 1203 gut microbiomes derived from faecal samples from the participants was also performed. Several associations between gut microbes and specific nutrients, foods, food groups and general dietary indices were found, which were driven by the presence and diversity of healthy and plant-based foods [5]. Overall microbiome composition was predictive for many cardiometabolic blood markers including fasting and postprandial glycaemic, lipidemic and inflammatory indices. Microbiome signatures grouped both microbiome and dietary components into health-associated clusters that were in agreement with dietary quality and diversity scores.

A Dutch study showed that dietary patterns derived from cluster analyses of food frequency questionnaires (FFQs), associated with pro-inflammatory and anti-inflammatory features of the gut microbiome [6]. Shotgun metagenomics sequencing was performed in faecal samples from 1425 individuals to investigate gut microbial composition and function. Processed foods and animal-derived foods were consistently associated with higher abundances of Firmicutes, *Ruminococcus* species of the *Blautia* genus and endotoxin synthesis pathways. On the other hand, plant foods and fish correlated positively with SCFA-producing bacteria and pathways of nutrient metabolism. Gut bacteria known for their shared function in health and disease were consistently associated with the identified dietary patterns. In addition, specific foods and nutrients correlated with bacterial species that are known to have anti-inflammatory effects and protective effects on the gut mucosa. These diet–gut microbiome associations were found both in patients with intestinal disease and the general population [6].

### Effect of diet on gut microbiota and cardiometabolic risk factors

It is primarily fibres from plant-based foods that can be degraded to SCFA by intestinal bacteria. Mice fed a diet without soluble fibre developed inflammation in the gut and poor intestinal health, which in turn led to weight gain [47]. Intestinal health was restored after soluble fibre was reintroduced into the diet. Several mice studies also show that high-fat diets, in particular high-fat diets rich in long-chain saturated fatty acids (SFA), are linked to unfavourable changes in type and numbers of gut



bacteria, resulting in dysbiosis and inflammation, with a subsequent increased risk of chronic disease, such as obesity and metabolic syndrome [48–51]. Dietary intervention trials in humans suggest that the microbiota-mediated effect of a dietary change on metabolism and health may be modest [52–55].

#### Human studies with high-fat diets

In people who ate a very low-carbohydrate diet which contained little carbohydrate (4 E %) and fibre and correspondingly more fat (61 E %) and protein (35 E %), a lower bacterial colonization in the colon has been shown [56]. Faecal butyrate concentrations and abundance of the *Roseburia/E. rectale* group are both reduced in people eating a carbohydrate-reduced diet (24 g/day) [57]. In a randomised controlled-feeding trial Wan et al. [52] compared three dietary patterns differing in carbohydrate and fat proportions: a lower-fat diet (fat 20% energy), a moderate-fat diet (fat 30% energy) and a higher-fat diet (fat 40% energy). The study was performed among 217 young, healthy adults during a 6 month's period. The researchers showed that the high-fat diet had unfavourable effects on the gut microbiota, faecal bacterial metabolites, and markers of inflammation, whereas the lower-fat diet was associated with a more favourable profile of these biomarkers.

In a recent systematic review, both interventional and observational studies showed associations between high fat diet intake, mainly rich in SFA, and reduction of bacterial abundance, diversity, and richness in the gut [4]. The dietary intervention studies showed no strong effects on gut microbiota and no association with metabolic outcomes. However, in observational studies high intake of total fat was positively correlated with the abundance of *Clostridium bolteae* and circulating serum levels of SFA correlated with *Blautia*, and both bacteria associated with unhealthy metabolic outcomes, i.e., insulin resistance and higher BMI or waist circumference [4]. Results from studies on diets rich in monounsaturated fatty acids (MUFA) were less consistent with no or possibly negative effects on total bacterial numbers and gut microbiota richness and diversity. In contrast, diets rich in omega-3 or omega-6 polyunsaturated fatty acids (PUFA) did not seem to affect the gut microbiota or metabolic health outcomes negatively. PUFA-enriched diets were associated with increased abundance of the Tenericutes phylum which in turn was associated with lower levels of triglycerides in plasma [4].

Evidence from randomised trials assessing the effect of PUFA on human gut microbiota is scarce. In a randomised cross-over trial Watson et al. [58] showed that a daily intake of 4 g omega-3 PUFA supplement (administered in capsules or drinks) over 8 weeks in 22

middle-aged, healthy volunteers was associated with reversible changes at gut family and genus levels, including an increase in the SCFA producing *Bifidobacterium*, *Lactobacillus*, *Lachnospira* and *Roseburia*. The authors concluded that the increase in density of butyrate producers was in line with existing preclinical literature and compatible with the known anti-inflammatory properties of omega-3 PUFA.

#### Human studies with high-fibre diets

Intervention trials in humans with fibre supplements or fibre enriched diets have consistently shown a positive effect on gut microbiota composition, with an increase in SCFA-producing bacteria and SCFA in faeces or blood samples [59–63]. Wheat bran supplementation (>70% arabinoxylan oligo-saccharides) increased the abundance of butyrate, acetate, and propionate as well as total SCFA concentrations in a human intervention trial [61]. However, increased faecal bulking and reduced transit time seen with increased dietary fibre, would decrease colonic absorption of SCFA and could partly explain the increases in faecal SCFA concentrations observed in studies with increased dietary fibre content [64].

A systematic review and meta-analysis of the effect of dietary interventions with fibre (mainly supplements) on gut microbiota in people with type 2 diabetes, showed that dietary fibre improved the relative abundance of *Bifidobacterium* and total SCFA, and improved glycosylated haemoglobin [65]. This systematic review included our own intervention study with 16 g per day of ITF for 6 weeks which induced moderate alterations in the composition of faecal bacteria, with an increased concentration of bifidobacteria being the most pronounced effect [66]. Compared to placebo, the prebiotic treatment also increased faecal concentrations of total SCFA, acetate, and propionate, but did not positively affect butyrate or the overall bacterial diversity [66]. Furthermore, the prebiotics had no positive effect on concentrations of glucose, insulin, gut hormones (GLP-1, GLP-2, PYY and ghrelin), appetite ratings or energy intake [67, 68].

Evidence from a recent systematic review suggests that ITF have a prebiotic effect on the gut microbiota, promoting the abundances of *Bifidobacterium*, *Lactobacillus*, and *Faecalibacterium prausnitzii* [69]. Beneficial health effects reported after intake of ITF included improved intestinal barrier function and laxation, increased insulin sensitivity, improved lipid profile, increased absorption of calcium and magnesium, and increased satiety [69]. However, another recent systematic review of ITF interventions in humans observed favourable effects of ITF intake on blood glucose, total cholesterol, and triglyceride concentration only in subjects with prediabetes and diabetes [70].

Three days with an evening meal of barley-kernel based bread rich in  $\beta$ -glucan fibres, increased fermentation activity and serum levels of SCFA in healthy adults, resulting in increased levels of gut hormones involved in regulation of blood glucose and appetite (GLP-1, PYY, and GLP-2), as well as improved insulin sensitivity [71]. Another study showed that a supplement with 3 g/d high molecular weight  $\beta$ -glucan increased Bacteroidetes and decreased Firmicutes abundances compared to placebo. At the genus level, the  $\beta$ -glucan supplement increased *Bacteroides*, and tended to increase *Prevotella* while decreasing *Dorea* [72]. However, low molecular weight  $\beta$ -glucan did not alter the intestinal microbiota composition [72]. The abundance of *Bacteroides*, *Prevotella*, and *Dorea* correlated with changes in risk factors for cardiovascular disease, such as BMI, waist circumference, blood pressure, and triglyceride levels [72]. This indicates that high molecular weight  $\beta$ -glucan can induce shifts in the intestinal microbiota that may partly explain the beneficial effect of  $\beta$ -glucan fibres on metabolism [72].

### SCFA and effects on metabolism

#### SCFA production, uptake, and turnover

Many of the beneficial health effects of dietary fibre, including prebiotic fibres are believed to be mediated through the microbial production of SCFA [73–76]. The SCFA mainly comprise acetate, propionate and butyrate, but include formate and lactate as well [75]. The molar ratio of acetate, propionate and butyrate in faeces is approximately 60:20:20 [77]. The bacteria can also metabolise lactate into acetate, propionate and butyrate [75].

SCFA are readily absorbed at similar rates in different parts of the colon [78]. They are then metabolized at three major sites in the body: ceco-colonic epithelium, liver cells and muscle cells [79]. Butyrate is the major energy source for ceco-colonic epithelium for maintenance-energy producing pathways [80]. Propionate is mainly used for gluconeogenesis in the liver, together with butyrate [81]. Acetate is also largely taken up by the liver. It enters the peripheral circulation to be metabolized by peripheral tissues, where oxidation of residual acetate is used for energy in muscle cells [82, 83]. Propionate and acetate appear less studied than butyrate, but may in addition to butyrate have anti-carcinogenic properties, and propionate is suggested to reduce visceral fat and liver fat [75].

SCFA act on host physiology through G protein-coupled receptors (GPCRs) and post-translational modifications [84, 85]. GPCRs are the largest receptor family in mammals, where six of them are sensitive to SCFA; GPR41 (also called free fatty acid receptor 3/FFAR3), GPR42, GPR43 (FFAR2), GPR109, GPR164, and olfactory receptor 78 (Olf78) [86, 87]. These receptors are found

in intestinal epithelium, immune cells, and fat cells, but their level of expression varies between tissues and cell types [85], and activation of the GPCRs may induce different effects in various tissues [88].

#### SCFA and inflammation

A large body of evidence suggests that an abnormal amount of lipids, hyperglycaemia and impaired insulin sensitivity can all cause endothelial dysfunction and low grade chronic inflammation, which have been linked to cardiovascular disease [13, 89, 90]. SCFA has a very potent anti-inflammatory effect, which block liberation of inflammatory mediators and, thus, reduces influx of immune cells to the site of inflammation, migration of immune cells, proliferation and persuades apoptosis [85, 91]. In this way functions mediated by GPCRs activation regulate the inflammatory process by preventing white blood cells from passing through the endothelium [92]. Furthermore, butyrate and acetate has been shown to exert beneficial effects on Angiotensin II-induced endothelial dysfunction in mice, by increasing the bio-accessibility of Nitric oxide (NO) and, thereby, reducing oxidative stress [93]. This effect was associated with GPR activation [93].

SCFA have effects on intestinal permeability and may, thus, regulate the systemic exposure to pro-inflammatory bacterial products, such as endotoxins or lipopolysaccharides (LPS) [94]. LPS are an essential part of the cell surface of most gram-negative bacteria [95, 96]. Translocation of LPS into the bloodstream induce development of low-grade endotoxaemia mostly by Toll-like receptor 4 (TLR4), acting as a receptor for LPS, signalling on macrophages, monocytes, and other cells of the inborn immune system. Furthermore, cluster of differentiation 14 (CD14) plays an important role in passing LPS to the TLR4 complex [97].

Patients with type 2 diabetes, overweight and atherosclerosis have increased levels of inflammatory cytokines, such as tumor necrosis factor alpha (TNF- $\alpha$ ) and interleucin-6 (IL-6) [98–101], and LPS, which is caused by low-grade endotoxaemia [97, 102, 103].

High-fat diets have been shown in studies with humans and mice to increase the degree of LPS-containing microbiota in the intestine [104–107]. Increased blood levels of LPS (endotoxaemia) are probably linked to changes in the intestinal microbiota, since antibiotic therapy reduces caecal and systemic LPS levels simultaneously with reduced glucose intolerance and fat mass development [97, 107, 108].

In an experiment in rats, a diet rich in fat and cholesterol enhanced the level of propionate simultaneously with a decline in butyrate, compared to a control group that received standard food for laboratory rats [109]. The

reduction in butyrate with time was linked to a raise in LPS liberated in the blood.

#### SCFA and regulation of appetite and blood glucose


SCFA have been shown to bind to GPR41 and GPR43 in enteroendocrine L-cells and thereby increase release of the gut hormones peptide YY (PYY), glucagon-like peptide-1 (GLP-1) and glucagon-like peptide-2 (GLP-2) after a meal (Fig. 4) [110]. This has the potential to improve glycaemic control and appetite regulation, as GLP-1 increases insulin release from pancreas and both GLP-1 and PYY are appetite suppressing hormones [111]. GLP-2 is important for preserving intestinal integrity [111, 112]. Animal studies show that butyrate improves glucose control by promoting gut production of GLP-1 and PYY as well as protecting the intestinal barrier [77, 113–115]. According to Sakakibara et al. [116], SCFA can also increase secretion of the appetite suppressing hormone leptin by activating GPR43, based on in vivo and in vitro studies.

#### SCFA and lipid metabolism

The SCFA can enter the circulation as substrates for lipid and cholesterol synthesis in the liver but can also be a regulatory factor in lipid metabolism [84]. SCFA

can enhance fatty acid oxidation and production of heat, block fatty acid synthesis, and reduce storage of fat in the body [117]. Angiopoietinlike 4 (ANGPTL4) is a signalling protein with several different functions that is synthesized in most tissues [118]. Studies suggest that ANGPTL4 is a key host protein that is reactive to the intestinal microbial environment. By controlling fatty acid uptake and metabolism in the tissues, ANGPTL4 can modify obesity in humans [119, 120]. SCFA, especially butyrate, affects lipid metabolism by inducing secretion of ANGPTL4 in human colon cell lines, which stimulates peroxisome proliferator-activated receptor- $\gamma$  (PPAR $\gamma$ ) and, thereby, blocks the activation of lipoprotein lipase (LPL) [119, 121]. LPL is important for the transfer of fatty acids from chylomicrons and very-low-density lipoprotein (VLDL) to adipocytes [122]. With low LPL activity, less fat is stored in adipose tissue [104, 123], and triglyceride-rich lipoproteins remain in the blood stream for a longer period, raising fat accumulation in and on the artery walls which may lead to atherosclerosis [124]. Furthermore, the excessive expression of ANGPTL4 in white adipose tissue decreases fat mass [125].

Finally, butyrate increases heat production and lipids utilisation through the uncoupling protein (UCP), which performs an essential role in lipid metabolism, and also



**Fig. 4** Effect of dietary fibre on glycaemic control and appetite, and possible pathways. **A** Dietary fibres escape digestion in the small intestine and **B** are fermented into SCFA acids by gut bacteria in the colon. **C** The SCFA bind to G-protein coupled receptors in enteroendocrine L-cells. **D** This causes increased secretion of GLP-1, GLP-2, leptin and PYY in response to a meal. **E** GLP-1 improves regulation of blood glucose by enhancing release of insulin and suppressing release of glucagon from pancreas. GLP-1 also protects the beta-cells. GLP-1, leptin, and PYY enhance satiety by affecting the brain and the gastrointestinal system. GLP-2 maintains the intestinal barrier and may, thus, prevent systemic inflammation. GLP-1 and 2, glucagon-like peptide 1 and 2; PYY, peptide YY; SCFA, short-chain fatty acids. *Figure was produced using Servier Medical Art and reproduced from first author's thesis*

improves lipid metabolism by activating adiponectin [84].

However, the role of SCFA in lipid metabolism and obesity remains controversial. Turnbaugh et al. [18], demonstrated in 2006 that SCFA can result in additional weight gain due to contribution of extra calories in the obese mice. Perry et al. [126] also indicated in 2016 that increased acetate turnover led to increased obesity and impaired insulin sensitivity in rodents.

#### **SCFA and blood pressure regulation**

The three main SCFA made by the microbiota in the intestinal lumen regulate blood pressure through Olfr78 and GPR41 [127, 128]. Olfr78 knock-out mice are hypotensive [129], whereas GPR41 knock-out mice are hypertensive [130], suggesting that these pathways may be important in linking SCFA and host blood pressure control. Acetate and propionate operate via a complex interaction that results in renin secretion mediated through Olfr78 and counter-regulation through GPR41. Butyrate works via attenuation of angiotensin II-induced expression of renal prorenin receptors and renin [76]. Olfr78 is expressed at high rates in the renal juxtaglomerular apparatus, where it causes increased renin secretion in response to SCFA binding [129]. Furthermore, both Olfr78 and GPR41 are expressed in smooth muscle cells of small resistance vessels, where they differentially mediate vascular tone [129].

#### **Human studies with SCFA supplementation**

Human studies with direct supplementation with SCFA are limited but largely in line with findings from animal models and cell-lines that support a beneficial role of SCFA in regulation of body weight, appetite, and energy expenditure as well as glycaemic control and insulin sensitivity [59]. In a study by Chambers et al. [131], propionate supplementation seemed to protect against weight gain when it was given to people as part of a habitual diet. Three other studies in humans have also shown that supplementation with propionate, targeting delivery in the lower gastrointestinal tract, may reduce energy intake [131–133]. Furthermore, human studies also support the observation in rodents that SCFA stimulate whole-body lipid oxidation and thereby increases energy expenditure [134–136]. Blaak et al. [59] summarise the findings from these and other studies in a recent review, where they conclude that SCFA administration studies and dietary intervention studies with prebiotics with the aim to increase SCFA production in humans, provides direct and indirect evidence for a beneficial effect of SCFA on blood glucose regulation and insulin sensitivity. Nevertheless, this has not been shown in all recent well-controlled studies, and Blaak et al. [59] mention that the lack

of effect is mostly observed in metabolically disturbed phenotypes, which suggests a disturbed SCFA handling/signalling in these individuals [59]. Interestingly, this is in accordance with our findings of no effect of ITF on glycaemic control and appetite regulation in patients with type 2 diabetes [67, 68], but is in contrast to the findings of Liu et al. [70] mentioned earlier, who found that interventions with ITF only had favourable effects on metabolic outcomes in people with prediabetes and diabetes.

#### **Conclusions and future perspectives**

In this narrative review, we have summarised research supporting the important role of diet in shaping gut microbiota and the intestinal production of SCFA. SCFA seem to have a significant impact on metabolic regulation and can, therefore, modulate cardiometabolic risk. This research field is still relatively young, so robust conclusions about causality cannot be made. Even though large methodological advances have been made, allowing us to investigate both the composition and the functionalities of the gut microbiota, limitations still exist in studying interaction between diet and gut microbiota in humans. Animal and cell line studies have paved the way for hypotheses to be explored in human studies, and the limited evidence we have from human studies so far, seem mainly to support the beneficial role of SCFA in regulation of blood glucose, blood lipids, and energy homeostasis. However, the underlying mechanisms for SCFA effects are still just being uncovered, and these studies are difficult to replicate in humans.

To broaden our understanding of diet-microbiota interaction in cardiometabolic health, and the role of SCFA in this interplay, several knowledge gaps have to be filled and methodology improved: First, we need better and more detailed and precise dietary investigation methods. The current investigation methods, like FFQs, are inherently imprecise. The use of dietary records, biomarkers of dietary intake and standardized meal challenges can complement information from FFQs. It is important to look beyond nutrients and single foods in diet-microbiome research, and one way of achieving a better understanding of this interaction is to perform unsupervised clustering analysis to identify dietary patterns and microbial clusters, like they did in the study by Bolte et al. [6].

Second, we should strive to develop more detailed and precise methods to identify and classify microbiota and SCFA in the human gut. Today's analyses in human studies are mainly restricted to faecal analyses whereas the main part of the SCFA is absorbed and used as substrate in the colonocytes. As just a minor percentage of the SCFA produced in the gut are excreted in the faeces, faecal analyses of SCFA may give a false impression of



dietary interventions' effect on gut microbiota and SCFA production.

Finally, we need a better understanding of the relationship between gut bacteria and human health. In the PREDICT study, the strongest microbiome–habitual diet associations were driven by poorly characterized microorganisms, which strongly suggests that our knowledge of the bacteria and their metabolic functions, is still far from complete and needs further investigation [5]. Future investigations of the remaining microbes that accompany the bacteria in the gut may contribute to higher understanding of human health and means to conquer illness. Further research, especially large, well-powered, long-term human intervention studies, is required to further understand and promote the role diet plays in modulating the gut microbiota and SCFA production and, thereby, human metabolism and health. These studies should be performed in populations that differs both geographically and with regard to sex, age, metabolic phenotypes, and health.

This narrative review gives an updated overview of a rapidly evolving research field that includes many different research disciplines and tries to connect them to yield a deeper understanding. At the same time, such a multidisciplinary approach may overlook details pertaining to the individual disciplines. Importantly, the clinical bearing of experimental basic studies must eventually be tested in well-designed clinical trials. The greater part of the research in this field is in-vitro and animal-studies. We chose to focus on human studies in this review and may thus have left out other types of studies of interest. Many of the human intervention studies included are acute studies and most of the long-term studies had a relative short duration of three to 12 weeks and a limited number of participants ( $n=6-30$ ) [4, 59, 63]. Our findings are also limited by the fact that the literature was gathered in a non-systematic way which may have caused some selection bias, as well as the fact that we did not formally analyse risk of bias in the intervention studies included.

What studies have found to promote diversity, increase SCFA production and be associated with beneficial effects on cardiometabolic risk factors, seems to be in line with the general guidelines for a healthy diet with an emphasize on plant-based foods like vegetables, fruit, berries, whole grains, nuts and fish. Furthermore, an unfavourable cardiometabolic profile is associated with highly processed foods rich in sugars, refined carbohydrates, fat, and saturated fat in particular. However, the metabolic responses to dietary interventions seem to vary depending on individual traits such as sex and existing gut microbiota, as well as genetics. With increasing knowledge about factors shaping the gut microbiota, the

potential to develop personalized dietary recommendations in prevention and treatment of disease and metabolic disturbances seems promising.

#### Abbreviations

ANGPTL4	Angiopoietin-like 4
FFAR	Free fatty acid receptor
CD14	Cluster of differentiation 14
FFQ	Food frequency questionnaire
GLP-1	Glucagon-like peptide-1
GLP-2	Glucagon-like peptide-2
GOS	Galacto-oligosaccharides
GPCR	G protein-coupled receptor
IL-6	Interleukin 6
ITF	Inulin-type fructans
LPL	Lipoprotein lipase
LPS	Lipopolysaccharides
MUFA	Monounsaturated fatty acids
Olf78	Olfactory receptor 78
PYY	Peptide YY
PUFA	Polyunsaturated fatty acids
SCFA	Short-chain fatty acids
SFA	Saturated fatty acids
TLR	Toll-like receptor
TNF- $\alpha$	Tumor necrosis factor alpha
TMA	Trimethylamine
UCP	Uncoupling protein

#### Authors' contributions

AMA, EB and SG performed the relevant background and literature searches for the construction of this review. All authors contributed in writing and editing, and read and approved the final manuscript.

#### Funding

Open access funding provided by University of Oslo (incl Oslo University Hospital)

#### Availability of data and materials

Not applicable.

#### Declarations

#### Ethics approval and consent to participate

Not applicable.

#### Consent for publication

All authors consent to this publication.

#### Competing interests

The authors declare no competing interests.

Received: 11 August 2022 Accepted: 7 March 2023

Published online: 14 March 2023

#### References

1. Afshin A, Forouzanfar MH, Reitsma MB, Sur P, Estep K, Lee A, et al. Health effects of overweight and obesity in 195 countries over 25 years. *N Engl J Med*. 2017;377(1):13–27.
2. Davies MJ, Aroda VR, Collins BS, Gabbay RA, Green J, Maruthur NM, et al. Management of hyperglycemia in type 2 diabetes, 2022 a consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care*. 2022;45(11):2753–86.
3. Lichtenstein AH, Appel LJ, Vadiveloo M, Hu FB, Kris-Etherton PM, Rebholz CM, et al. 2021 dietary guidance to improve cardiovascular health:

- a scientific statement from the American Heart Association. *Circulation*. 2021;144(23):e472–87.
4. Wolters M, Ahrens J, Romani-Pérez M, Watkins C, Sanz Y, Benítez-Páez A, et al. Dietary fat, the gut microbiota, and metabolic health - A systematic review conducted within the MyNewGut project. *Clinical Nutrition* (Edinburgh, Scotland). 2019;38(6):2504–20.
  5. Asnicar F, Berry SE, Valdes AM, Nguyen LH, Piccinno G, Drew DA, et al. Microbiome connections with host metabolism and habitual diet from 1,098 deeply phenotyped individuals. *Nat Med*. 2021;27:321.
  6. Bolte LA, Vich Vila A, Imhann F, Collij V, Gacesa R, Peters V, et al. Long-term dietary patterns are associated with pro-inflammatory and anti-inflammatory features of the gut microbiome. *Gut*. 2021;70(7):1287–98.
  7. Zeevi D, Korem T, Zmora N, Israeli D, Rothschild D, Weinberger A, et al. Personalized nutrition by prediction of glycemic responses. *Cell*. 2015;163(5):1079–94.
  8. Qin J, Li Y, Cai Z, Li S, Zhu J, Zhang F, et al. A metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature*. 2012;490(7418):55–60.
  9. Le Chatelier E, Nielsen T, Qin J, Prifti E, Hildebrand F, Falony G, et al. Richness of human gut microbiome correlates with metabolic markers. *Nature*. 2013;500(7464):541–6.
  10. Lepage P, Häslér R, Spehlmann ME, Rehman A, Zvirbliene A, Begun A, et al. Twin study indicates loss of interaction between microbiota and mucosa of patients with ulcerative colitis. *Gastroenterology*. 2011;141(1):227–36.
  11. Manichanh C, Borruel N, Casellas F, Guarnier F. The gut microbiota in IBD. *Nat Rev Gastroenterol Hepatol*. 2012;9(10):599–608.
  12. Karlsson FH, Tremaroli V, Nookaew I, Bergstrom G, Behre CJ, Fagerberg B, et al. Gut metagenome in European women with normal, impaired and diabetic glucose control. *Nature*. 2013;498(7452):99–103.
  13. Peh A, O'Donnell JA, Broughton BRS, Marques FZ. Gut microbiota and their metabolites in stroke: a double-edged sword. *Stroke*. 2022;53(5):1788–801.
  14. Berry SE, Valdes AM, Drew DA, Asnicar F, Mazidi M, Wolf J, et al. Human postprandial responses to food and potential for precision nutrition. *Nat Med*. 2020;26(6):964–73.
  15. Ursell LK, Metcalf JL, Parfrey LW, Knight R. Defining the human microbiome. *Nutr Rev*. 2012;70(1):38–44.
  16. Mohajeri MH, Brummer RJM, Rastall RA, Weersma RK, Harmsen HJM, Faas M, et al. The role of the microbiome for human health: from basic science to clinical applications. *Eur J Nutr*. 2018;57:1.
  17. Bäckhed F, Ding H, Wang T, Hooper LV, Koh GY, Nagy A, et al. The gut microbiota as an environmental factor that regulates fat storage. *Proc Natl Acad Sci USA*. 2004;101(44):15718–23.
  18. Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JL. An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature*. 2006;444(7122):1027–31.
  19. Karlsson FH, Fåk F, Nookaew I, Tremaroli V, Fagerberg B, Petranovic D, et al. Symptomatic atherosclerosis is associated with an altered gut metagenome. *Nat Commun*. 2012;3:1245.
  20. Brüssow H. Problems with the concept of gut microbiota dysbiosis. *Microb Biotechnol*. 2020;13(2):423–34.
  21. Sender R, Fuchs S, Milo R. Revised estimates for the number of human and bacterial cells in the body. *PLoS Biol*. 2016;14(8):e1002533.
  22. Gilbert JA, Blaser MJ, Caporaso JG, Jansson JK, Lynch SV, Knight R. Current understanding of the human microbiome. *Nat Med*. 2018;24(4):392–400.
  23. Stanislawski MA, Dabelea D, Lange LA, Wagner BD, Lozupone CA. Gut microbiota phenotypes of obesity. *NPJ Biofilms Microbiomes*. 2019;5(1):18.
  24. Forslund K, Hildebrand F, Nielsen T, Falony G, Le Chatelier E, Sunagawa S, et al. Disentangling type 2 diabetes and metformin treatment signatures in the human gut microbiota. *Nature*. 2015;528(7581):262–6.
  25. Jackson MA, Verdi S, Maxan ME, Shin CM, Zierer J, Bowyer RCE, et al. Gut microbiota associations with common diseases and prescription medications in a population-based cohort. *Nat Commun*. 2018;9(1):2655.
  26. Rothschild D, Weissbrod O, Barkan E, Kurilshikov A, Korem T, Zeevi D, et al. Environment dominates over host genetics in shaping human gut microbiota. *Nature*. 2018;555(7695):210–5.
  27. Ridaura VK, Faith JJ, Rey FE, Cheng J, Duncan AE, Kau AL, et al. Gut microbiota from twins discordant for obesity modulate metabolism in mice. *Science*. 2013;341(6150):1241214.
  28. Faith JJ, Guruge JL, Charbonneau M, Subramanian S, Seedorf H, Goodman AL, et al. The long-term stability of the human gut microbiota. *Science*. 2013;341(6141):1237439.
  29. Human Microbiome Project Consortium. Structure, function and diversity of the healthy human microbiome. *Nature*. 2012;486(7402):207–14.
  30. Lynch SV, Pedersen O. The human intestinal microbiome in health and disease. *N Engl J Med*. 2016;375(24):2369–79.
  31. Odamaki T, Kato K, Sugahara H, Hashikura N, Takahashi S, Xiao JZ, et al. Age-related changes in gut microbiota composition from newborn to centenarian: a cross-sectional study. *BMC Microbiol*. 2016;16:90.
  32. Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, et al. A human gut microbial gene catalogue established by metagenomic sequencing. *Nature*. 2010;464(7285):59–65.
  33. Gill SR, Pop M, Deboy RT, Eckburg PB, Turnbaugh PJ, Samuel BS, et al. Metagenomic analysis of the human distal gut microbiome. *Science*. 2006;312(5778):1355–9.
  34. Scientific Advisory Committee on Nutrition. Independent report commissioned by Department of Health and the Food Standards Agency. *Public Health England*. 2015. Carbohydrates and health. ISBN 978017082847 PDF, 2.39 MB, 384 pages.
  35. Flint HJ, Scott KP, Louis P, Duncan SH. The role of the gut microbiota in nutrition and health. *Nat Rev Gastroenterol Hepatol*. 2012;9(10):577–89.
  36. Fan Y, Pedersen O. Gut microbiota in human metabolic health and disease. *Nat Rev Microbiol*. 2021;19(1):55–71.
  37. Gibson GR, Hutkins R, Sanders ME, Prescott SL, Reimer RA, Salminen SJ, et al. Expert consensus document: The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nat Rev Gastroenterol Hepatol*. 2017;14(8):491–502.
  38. Kellow NJ, Coughlan MT, Reid CM. Metabolic benefits of dietary prebiotics in human subjects: a systematic review of randomised controlled trials. *Br J Nutr*. 2014;111(7):1147–61.
  39. Beserra BT, Fernandes R, do Rosario VA, Mocellin MC, Kuntz MG, Trindade EB. A systematic review and meta-analysis of the prebiotics and synbiotics effects on glycaemia, insulin concentrations and lipid parameters in adult patients with overweight or obesity. *Clinical Nutrition* (Edinburgh, Scotland). 2014;34:845.
  40. Bonsu NK, Johnson CS, McLeod KM. Can dietary fructans lower serum glucose? *J Diabetes*. 2011;3(1):58–66.
  41. da Silva ST, dos Santos CA, Bressan J. Intestinal microbiota; relevance to obesity and modulation by prebiotics and probiotics. *Nutr Hosp*. 2013;28(4):1039–48.
  42. Mahboobi S, Rahimi F, Jafarnejad S. Effects of prebiotic and synbiotic supplementation on glycaemia and lipid profile in type 2 diabetes: a meta-analysis of randomized controlled trials. *Adv Pharm Bull*. 2018;8(4):565–74.
  43. Liber A, Szajewska H. Effects of inulin-type fructans on appetite, energy intake, and body weight in children and adults: systematic review of randomized controlled trials. *Ann Nutr Metab*. 2013;63(1–2):42–54.
  44. John GK, Wang L, Nanavati J, Twose C, Singh R, Mullin G. Dietary alteration of the gut microbiome and its impact on weight and fat mass: a systematic review and meta-analysis. *Genes* (Basel). 2018;9(3):167.
  45. da Silva Borges D, Fernandes R, Thives Mello A, da Silva Fontoura E, Soares Dos Santos AR, Santos de Moraes Trindade EB. Prebiotics may reduce serum concentrations of C-reactive protein and ghrelin in overweight and obese adults: a systematic review and meta-analysis. *Nutr Rev*. 2020;78(3):235–48.
  46. Rao M, Gao C, Xu L, Jiang L, Zhu J, Chen G, et al. Effect of inulin-type carbohydrates on insulin resistance in patients with type 2 diabetes and obesity: a systematic review and meta-analysis. *J Diabetes Res*. 2019;2019:5101423.
  47. Chassaing B, Miles-Brown J, Pellizzon M, Ulman E, Ricci M, Zhang L, et al. Lack of soluble fiber drives diet-induced adiposity in mice. *Am J Physiol Gastrointest Liver Physiol*. 2015;309(7):G528–41.
  48. Cani PD, Neyrinck AM, Fava F, Knauf C, Burcelin RG, Tuohy KM, et al. Selective increases of bifidobacteria in gut microflora improve high-fat-diet-induced diabetes in mice through a mechanism associated with endotoxaemia. *Diabetologia*. 2007;50(11):2374–83.

49. Pendyala S, Walker JM, Holt PR. A high-fat diet is associated with endotoxemia that originates from the gut. *Gastroenterology*. 2012;142(5):1100–1.e2.
50. Caesar R, Tremaroli V, Kovatcheva-Datchary P, Cani PD, Bäckhed F. Crosstalk between gut microbiota and dietary lipids aggravates WAT inflammation through TLR signaling. *Cell Metab*. 2015;22(4):658–68.
51. Lam YY, Ha CW, Hoffmann JM, Oscarsson J, Dinudom A, Mather TJ, et al. Effects of dietary fat profile on gut permeability and microbiota and their relationships with metabolic changes in mice. *Obesity (Silver Spring, Md)*. 2015;23(7):1429–39.
52. Wan Y, Wang F, Yuan J, Li J, Jiang D, Zhang J, et al. Effects of dietary fat on gut microbiota and faecal metabolites, and their relationship with cardiometabolic risk factors: a 6-month randomised controlled-feeding trial. *Gut*. 2019;68(8):1417–29.
53. Wu GD, Chen J, Hoffmann C, Bittinger K, Chen YY, Keilbaugh SA, et al. Linking long-term dietary patterns with gut microbial enterotypes. *Science*. 2011;334(6052):105–8.
54. Cotillard A, Kennedy SP, Kong LC, Prifti E, Pons N, Le Chatelier E, et al. Dietary intervention impact on gut microbial gene richness. *Nature*. 2013;500(7464):585–8.
55. Wu GD, Compher C, Chen EZ, Smith SA, Shah RD, Bittinger K, et al. Comparative metabolomics in vegans and omnivores reveal constraints on diet-dependent gut microbiota metabolite production. *Gut*. 2016;65(1):63–72.
56. Brinkworth GD, Noakes M, Clifton PM, Bird AR. Comparative effects of very low-carbohydrate, high-fat and high-carbohydrate, low-fat weight-loss diets on bowel habit and faecal short-chain fatty acids and bacterial populations. *Br J Nutr*. 2009;101(10):1493–502.
57. Duncan SH, Belonguer A, Holtrop G, Johnstone AM, Flint HJ, Lobley GE. Reduced dietary intake of carbohydrates by obese subjects results in decreased concentrations of butyrate and butyrate-producing bacteria in feces. *Appl Environ Microbiol*. 2007;73(4):1073–8.
58. Watson H, Mitra S, Croden FC, Taylor M, Wood HM, Perry SL, et al. A randomised trial of the effect of omega-3 polyunsaturated fatty acid supplements on the human intestinal microbiota. *Gut*. 2018;67(11):1974–83.
59. Blaak EE, Canfora EE, Theis S, Frost G, Groen AK, Mithieux G, et al. Short chain fatty acids in human gut and metabolic health. *Benef Microbes*. 2020;11(5):411–55.
60. Zhao L, Zhang F, Ding X, Wu G, Lam YY, Wang X, et al. Gut bacteria selectively promoted by dietary fibers alleviate type 2 diabetes. *Science*. 2018;359(6380):1151–6.
61. François IE, Lescroart O, Veraverbeke WS, Marzorati M, Possemiers S, Evenepoel P, et al. Effects of a wheat bran extract containing arabinoxylan oligosaccharides on gastrointestinal health parameters in healthy adult human volunteers: a double-blind, randomised, placebo-controlled, cross-over trial. *Br J Nutr*. 2012;108(12):2229–42.
62. O'Keefe SJ, Li JV, Lahti L, Ou J, Carbonero F, Mohammed K, et al. Fat, fibre and cancer risk in African Americans and rural Africans. *Nat Commun*. 2015;6:6342.
63. Swanson KS, de Vos WM, Martens EC, Gilbert JA, Menon RS, Soto-Vaca A, et al. Effect of fructans, prebiotics and fibres on the human gut microbiome assessed by 16S rRNA-based approaches: a review. *Benef Microbes*. 2020;11(2):101–29.
64. Rowland I, Gibson G, Heinken A, Scott K, Swann J, Thiele I, et al. Gut microbiota functions: metabolism of nutrients and other food components. *Eur J Nutr*. 2018;57(1):1–24.
65. Ojo O, Ojo OO, Zand N, Wang X. The effect of dietary fibre on gut microbiota, lipid profile, and inflammatory markers in patients with type 2 diabetes: a systematic review and meta-analysis of randomised controlled trials. *Nutrients*. 2021;13(6):1805.
66. Birkeland E, Gharagozian S, Birkeland KI, Valeur J, Måge I, Rud I, et al. Prebiotic effect of inulin-type fructans on faecal microbiota and short-chain fatty acids in type 2 diabetes: a randomised controlled trial. *Eur J Nutr*. 2020;59:3329.
67. Birkeland E, Gharagozian S, Birkeland KI, Holm OKS, Thorsby PM, Aas AM. Effect of inulin-type fructans on appetite in patients with type 2 diabetes: a randomised controlled crossover trial. *J Nutr Sci*. 2021;10:e72.
68. Birkeland E, Gharagozian S, Gulseth HL, Birkeland KI, Hartmann B, Holst JJ, et al. Effects of prebiotics on postprandial GLP-1, GLP-2 and glucose regulation in patients with type 2 diabetes: a randomised, double-blind, placebo-controlled crossover trial. *Diabet Med*. 2021;38:e14657.
69. Hughes RL, Alvarado DA, Swanson KS, Holscher HD. The prebiotic potential of inulin-type Fructans: a systematic review. *Adv Nutr*. 2021;13(2):492–529.
70. Li L, Li P, Xu L. Assessing the effects of inulin-type fructan intake on body weight, blood glucose, and lipid profile: A systematic review and meta-analysis of randomized controlled trials. *Food Sci Nutr*. 2021;9(8):4598–616.
71. Nilsson AC, Johansson-Boll EV, Björck IM. Increased gut hormones and insulin sensitivity index following a 3-d intervention with a barley kernel-based product: a randomised cross-over study in healthy middle-aged subjects. *Br J Nutr*. 2015;114(6):899–907.
72. Wang Y, Ames NP, Tun HM, Tosh SM, Jones PJ, Khafipour E. High molecular weight Barley  $\beta$ -Glucan alters gut microbiota toward reduced cardiovascular disease risk. *Front Microbiol*. 2016;7:129.
73. Correa-Oliveira R, Fachi JL, Vieira A, Sato FT, Vinolo MA. Regulation of immune cell function by short-chain fatty acids. *Clin Transl Immunol*. 2016;5(4):e73.
74. Donohoe DR, Garge N, Zhang X, Sun W, O'Connell TM, Bunker MK, et al. The microbiome and butyrate regulate energy metabolism and autophagy in the mammalian colon. *Cell Metab*. 2011;13(5):517–26.
75. Morrison DJ, Preston T. Formation of short chain fatty acids by the gut microbiota and their impact on human metabolism. *Gut microbes*. 2016;7(3):189–200.
76. Chambers ES, Preston T, Frost G, Morrison DJ. Role of gut microbiota-generated short-chain fatty acids in metabolic and cardiovascular health. *Curr Nutr Rep*. 2018;7(4):198–206.
77. Hamer HM, Jonkers D, Venema K, Vanhoutvin S, Troost FJ, Brummer RJ. Review article: the role of butyrate on colonic function. *Aliment Pharmacol Ther*. 2008;22(2):104–19.
78. den Besten G, van Eunen K, Groen AK, Venema K, Reijngoud DJ, Bakker BM. The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *J Lipid Res*. 2013;54(9):2325–40.
79. Wong JM, de Souza R, Kendall CW, Emam A, Jenkins DJ. Colonic health: fermentation and short chain fatty acids. *J Clin Gastroenterol*. 2006;40(3):235–43.
80. De Vadder F, Kovatcheva-Datchary P, Goncalves D, Vinera J, Zitoun C, Duchamp A, et al. Microbiota-generated metabolites promote metabolic benefits via gut-brain neural circuits. *Cell*. 2014;156(1–2):84–96.
81. Puchowicz MA, Bederman IR, Comte B, Yang D, David F, Stone E, et al. Zonation of acetate labeling across the liver: implications for studies of lipogenesis by MIDA. *Am J Physiol*. 1999;277(6):E1022–7.
82. Cummings JH, Pomare EW, Branch WJ, Naylor CP, Macfarlane GT. Short chain fatty acids in human large intestine, portal, hepatic and venous blood. *Gut*. 1987;28(10):1221–7.
83. Hijova E, Chmelarova A. Short chain fatty acids and colonic health. *Bratisl Lek Listy*. 2007;108(8):354–8.
84. He J, Zhang P, Shen L, Niu L, Tan Y, Chen L, et al. Short-chain fatty acids and their association with signalling pathways in inflammation, glucose and lipid metabolism. *Int J Mol Sci*. 2020;21(17):6356.
85. Ohira H, Tsutsui W, Fujioka Y. Are short chain fatty acids in gut microbiota defensive players for inflammation and atherosclerosis? *J Atheroscler Thromb*. 2017;24(7):660–72.
86. Canals M, Poole DP, Veldhuis NA, Schmidt BL, Bunnett NW. G-protein-coupled receptors are dynamic regulators of digestion and targets for digestive diseases. *Gastroenterology*. 2019;156(6):1600–16.
87. Suchý T, Zieschang C, Popkova Y, Kaczmarek I, Weiner J, Liebing AD, et al. The repertoire of adhesion G protein-coupled receptors in adipocytes and their functional relevance. *Int J Obes*. 2020;44(10):2124–36.
88. Brown AJ, Goldsworthy SM, Barnes AA, Eilert MM, Tcheang L, Daniels D, et al. The orphan G protein-coupled receptors GPR41 and GPR43 are activated by propionate and other short chain carboxylic acids. *J Biol Chem*. 2003;278(13):11312–9.
89. Grundy SM, Cleeman JJ, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and management of the metabolic syndrome: an American heart association/national heart, lung, and blood institute scientific statement. *Circulation*. 2005;112(17):2735–52.
90. Ades PA, Savage PD. Potential benefits of weight loss in coronary heart disease. *Prog Cardiovasc Dis*. 2014;56(4):448–56.

91. Nogal A, Valdes AM, Menni C. The role of short-chain fatty acids in the interplay between gut microbiota and diet in cardio-metabolic health. *Gut Microbes*. 2021;13(1):1–24.
92. González-Bosch C, Boorman E, Zunszain PA, Mann GE. Short-chain fatty acids as modulators of redox signaling in health and disease. *Redox Biol*. 2021;47:102165.
93. Robles-Vera I, Toral M, de la Visitación N, Aguilera-Sánchez N, Redondo JM, Duarte J. Protective effects of short-chain fatty acids on endothelial dysfunction induced by angiotensin II. *Front Physiol*. 2020;11:277.
94. Candelli M, Franza L, Pignataro G, Ojetti V, Covino M, Piccioni A, et al. Interaction between lipopolysaccharide and gut microbiota in inflammatory bowel diseases. *Int J Mol Sci*. 2021;22(12):6242.
95. Avila-Calderón ED, Ruiz-Palma MDS, Aguilera-Arreola MG, Velázquez-Guadarrama N, Ruiz EA, Gomez-Lunar Z, et al. Outer membrane vesicles of gram-negative bacteria: an outlook on biogenesis. *Front Microbiol*. 2021;12:557902.
96. Czepiel J, Biesiada G, Brzozowski T, Ptak-Belowska A, Perucki W, Birczynska M, et al. The role of local and systemic cytokines in patients infected with *Clostridium difficile*. *J Physiol Pharmacol J Polish Physiol Soc*. 2014;65(5):695–703.
97. Trøseid M, Nestvold TK, Rudi K, Thoresen H, Nielsen EW, Lapppegård KT. Plasma lipopolysaccharide is closely associated with glycemic control and abdominal obesity: evidence from bariatric surgery. *Diabetes Care*. 2013;36(11):3627–32.
98. Bakker GC, van Erk MJ, Pellis L, Wopereis S, Rubingh CM, Cnubben NH, et al. An antiinflammatory dietary mix modulates inflammation and oxidative and metabolic stress in overweight men: a nutrigenomics approach. *Am J Clin Nutr*. 2010;91(4):1044–59.
99. Krogh-Madsen R, Plomgaard P, Møller K, Mittendorfer B, Pedersen BK. Influence of TNF- $\alpha$  and IL-6 infusions on insulin sensitivity and expression of IL-18 in humans. *Am J Physiol Endocrinol Metab*. 2006;291(1):E108–14.
100. Esposito K, Nappo F, Marfella R, Giugliano G, Giugliano F, Ciotola M, et al. Inflammatory cytokine concentrations are acutely increased by hyperglycemia in humans: role of oxidative stress. *Circulation*. 2002;106(16):2067–72.
101. Kern PA, Saghizadeh M, Ong JM, Bosch RJ, Deem R, Simsolo RB. The expression of tumor necrosis factor in human adipose tissue. Regulation by obesity, weight loss, and relationship to lipoprotein lipase. *J Clin Invest*. 1995;95(5):2111–9.
102. Liang H, Hussey SE, Sanchez-Avila A, Tantiwong P, Musi N. Effect of lipopolysaccharide on inflammation and insulin action in human muscle. *PLoS ONE*. 2013;8(5):e63983.
103. Violi F, Cammisotto V, Bartimoccia S, Pignatelli P, Carnevale R, Nocella C. Gut-derived low-grade endotoxaemia, atherothrombosis and cardiovascular disease. *Nat Rev Cardiol*. 2023;20(1):24–37.
104. Cani PD, Delzenne NM. The role of the gut microbiota in energy metabolism and metabolic disease. *Curr Pharm Des*. 2009;15(13):1546–58.
105. Cani PD, Delzenne NM. Gut microflora as a target for energy and metabolic homeostasis. *Curr Opin Clin Nutr Metab Care*. 2007;10(6):729–34.
106. Delzenne NM, Cani PD. Interaction between obesity and the gut microbiota: relevance in nutrition. *Annu Rev Nutr*. 2011;31:15–31.
107. Cani PD, Bibiloni R, Knauf C, Waget A, Neyrinck AM, Delzenne NM, et al. Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity and diabetes in mice. *Diabetes*. 2008;57(6):1470–81.
108. Fujisaka S, Ussar S, Clish C, Devkota S, Dreyfuss JM, Sakaguchi M, et al. Antibiotic effects on gut microbiota and metabolism are host dependent. *J Clin Invest*. 2016;126(12):4430–43.
109. Maciejewska D, Skonieczna-Zydecka K, Lukomska A, Gutowska I, Dec K, Kupnicka P, et al. The short chain fatty acids and lipopolysaccharides status in Sprague-Dawley rats fed with high-fat and high-cholesterol diet. *J Physiol Pharmacol J Polish Physiol Soc*. 2018;69(2):205–10.
110. Holst JJ. The physiology of glucagon-like peptide 1. *Physiol Rev*. 2007;87(4):1409–39.
111. Cani PD, Everard A, Duparc T. Gut microbiota, enteroendocrine functions and metabolism. *Curr Opin Pharmacol*. 2013;13(6):935–40.
112. Hare KJ, Vilsboll T, Asmar M, Deacon CF, Knop FK, Holst JJ. The glucagonostatic and insulinotropic effects of glucagon-like peptide 1 contribute equally to its glucose-lowering action. *Diabetes*. 2010;59(7):1765–70.
113. Scheppach W, Weiler F. The butyrate story: old wine in new bottles? *Curr Opin Clin Nutr Metab Care*. 2004;7(5):563–7.
114. Lin HV, Frassetto A, Kowalik EJ Jr, Nawrocki AR, Lu MM, Kosinski JR, et al. Butyrate and propionate protect against diet-induced obesity and regulate gut hormones via free fatty acid receptor 3-independent mechanisms. *PLoS ONE*. 2012;7(4):e35240.
115. Gao Z, Yin J, Zhang J, Ward RE, Martin RJ, Lefevre M, et al. Butyrate improves insulin sensitivity and increases energy expenditure in mice. *Diabetes*. 2009;58(7):1509–17.
116. Sakakibara S, Yamauchi T, Oshima Y, Tsukamoto Y, Kadowaki T. Acetic acid activates hepatic AMPK and reduces hyperglycemia in diabetic KK-A(y) mice. *Biochem Biophys Res Commun*. 2006;344(2):597–604.
117. Kimura I, Ichimura A, Ohue-Kitano R, Igarashi M. Free fatty acid receptors in health and disease. *Physiol Rev*. 2020;100(1):171–210.
118. Fernández-Hernando C, Suárez Y. ANGPTL4: a multifunctional protein involved in metabolism and vascular homeostasis. *Curr Opin Hematol*. 2020;27(3):206–13.
119. Barja-Fernández S, Folgueira C, Castelao C, Pena-León V, González-Saenz P, Vázquez-Cobela R, et al. ANGPTL4 is associated with obesity and lipid profile in children and adolescents. *Nutrients*. 2019;11(6):1340.
120. Ortega-Senovilla H, van Poppel MNM, Desoye G, Herrera E. Angiopoietin-like protein 4 (ANGPTL4) is related to gestational weight gain in pregnant women with obesity. *Sci Rep*. 2018;8(1):12428.
121. Robciuc MR, Skrobuk P, Anisimov A, Oikkonen VM, Alitalo K, Eckel RH, et al. Angiopoietin-like 4 mediates PPAR delta effect on lipoprotein lipase-dependent fatty acid uptake but not on beta-oxidation in myotubes. *PLoS ONE*. 2012;7(10):e46212.
122. Alex S, Lange K, Amolo T, Grinstead JS, Haakonsson AK, Szalowska E, et al. Short-chain fatty acids stimulate angiopoietin-like 4 synthesis in human colon adenocarcinoma cells by activating peroxisome proliferator-activated receptor  $\gamma$ . *Mol Cell Biol*. 2013;33(7):1303–16.
123. Wellen KE, Hotamisligil GS. Inflammation, stress, and diabetes. *J Clin Invest*. 2005;115(5):1111–9.
124. Fantuzzi G, Mazzone T. Adipose tissue and atherosclerosis: exploring the connection. *Arterioscler Thromb Vasc Biol*. 2007;27(5):996–1003.
125. Delzenne NM, Neyrinck AM, Backhed F, Cani PD. Targeting gut microbiota in obesity: effects of prebiotics and probiotics. *Nat Rev Endocrinol*. 2011;7(11):639–46.
126. Perry RJ, Peng L, Barry NA, Cline GW, Zhang D, Cardone RL, et al. Acetate mediates a microbiome-brain- $\beta$ -cell axis to promote metabolic syndrome. *Nature*. 2016;534(7606):213–7.
127. Wu Y, Xu H, Tu X, Gao Z. The Role of Short-Chain Fatty Acids of Gut Microbiota Origin in Hypertension. *Front Microbiol*. 2021;12:730809.
128. Pluznick J. A novel SCFA receptor, the microbiota, and blood pressure regulation. *Gut microbes*. 2014;5(2):202–7.
129. Pluznick JL, Protzko RJ, Gevorgyan H, Peterlin Z, Sipos A, Han J, et al. Olfactory receptor responding to gut microbiota-derived signals plays a role in renin secretion and blood pressure regulation. *Proc Natl Acad Sci USA*. 2013;110(11):4410–5.
130. Natarajan N, Hori D, Flavahan S, Steppan J, Flavahan NA, Berkowitz DE, et al. Microbial short chain fatty acid metabolites lower blood pressure via endothelial G protein-coupled receptor 41. *Physiol Genomics*. 2016;48(11):826–34.
131. Chambers ES, Viardot A, Psichas A, Morrison DJ, Murphy KG, Zac-Varghese SE, et al. Effects of targeted delivery of propionate to the human colon on appetite regulation, body weight maintenance and adiposity in overweight adults. *Gut*. 2015;64(11):1744–54.
132. Byrne CS, Chambers ES, Alhabeed H, Chhina N, Morrison DJ, Preston T, et al. Increased colonic propionate reduces anticipatory reward responses in the human striatum to high-energy foods. *Am J Clin Nutr*. 2016;104(1):5–14.
133. Polyviou T, MacDougall K, Chambers ES, Viardot A, Psichas A, Jawaid S, et al. Randomised clinical study: inulin short-chain fatty acid esters for targeted delivery of short-chain fatty acids to the human colon. *Aliment Pharmacol Ther*. 2016;44(7):662–72.
134. Canfora EE, van der Beek CM, Jocken JWE, Goossens GH, Holst JJ, Olde Damink SWM, et al. Colonic infusions of short-chain fatty acid mixtures promote energy metabolism in overweight/obese men: a randomized crossover trial. *Sci Rep*. 2017;7(1):2360.
135. Chambers ES, Byrne CS, Aspey K, Chen Y, Khan S, Morrison DJ, et al. Acute oral sodium propionate supplementation raises resting energy



expenditure and lipid oxidation in fasted humans. *Diabetes Obes Metab.* 2018;20(4):1034–9.

136. van der Beek CM, Canfora EE, Lenaerts K, Troost FJ, Olde Damink SWM, Holst JJ, et al. Distal, not proximal, colonic acetate infusions promote fat oxidation and improve metabolic markers in overweight/obese men. *Clin Sci (Lond).* 2016;130(22):2073–82.

### **Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

